

## POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# INTERCEPT® 480SL

**ACTIVE CONSTITUENT: 480 g/L ETHEPHON**  
**(an anti-cholinesterase compound)**

A contraceptive treatment for the prevention  
of *Poa annua* seedhead emergence in golf greens

## 5 Litres

**PLANT PROTECTION**  
TURF GROWTH REGULATOR

Greenshed Pty Ltd ABN: 70 105 996 307  
2/17 Barclay St Marrickville NSW 2204  
Tel: 02 9667 4333 Fax: 02 9667 4999  
MATCHplayproducts.com.au

## GENERAL INSTRUCTIONS

### MIXING

Half fill the spray tank with water, add the required amount of product, then the remainder of water and mix well. DO NOT use alkaline water.

### EQUIPMENT

This product is corrosive to metals. Mix only the quantity of spray required for immediate use and do not allow to stand in metal containers for more than 2 hours. Flush equipment thoroughly with water after use.

### WARNING

DO NOT apply by hand held application methods.

### RE-ENTRY PERIOD:

#### PUBLIC

DO NOT allow public to access treated areas for 4 hours after application.

#### WORKERS

DO NOT allow entry into treated areas for 4 hours after application unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

### PROTECTION OF LIVESTOCK

This product is dangerous to bees and will kill bees which are foraging in the area to be treated.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that could be expected to cause drift onto nearby plants and adjacent crops, crop-lands or pasture. Wash sprayer thoroughly after use.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

## SAFETY DIRECTIONS

Harmful if inhaled. Corrosive. Attacks eyes and skin. Repeated minor exposure may have a cumulative poisoning effect. DO NOT inhale vapour or spray mist. Avoid contact with eyes and skin. When opening the container, preparing spray and using prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves, chemical resistant footwear and goggles.

If product or spray in eyes, wash it out immediately with water. If product or spray on skin, immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

## DIRECTIONS FOR USE

### RESTRAINTS

**DO NOT** re-apply INTERCEPT® 480SL within 2 weeks of the initial application.

**DO NOT** apply multiple applications of INTERCEPT® 480SL in areas where excessive thatch has accumulated.

**DO NOT** mix INTERCEPT® 480SL with Ammonium Thiosulphate.

**DO NOT** exceed a maximum of INTERCEPT® 480SL of 42 L/ha per year.

For use in all States where appropriate for the crop and/or pest.

CROP	PEST	RATE	CRITICAL COMMENTS
Gold Course Greens	Wintergrass ( <i>Poa annua</i> ) Seedhead Suppression	7-10 L/ha in 400-500 L/ha of water per hectare.	Make initial application prior to the emergence of new seedheads in Spring and Autumn. A period of 2-3 weeks after application is required for maximum seed head suppression performance. A repeat application is required in 14-21 days. Tank Mixtures of INTERCEPT® 480SL and PRIMO MAXX at 400 mL/ha can be used to promote seedhead suppression as well as turfgrass quality. Application of this mixture during frost periods may cause temporary turf discolouration.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD: DO NOT** graze treated turf/lawn; or feed turf/lawn clippings from any treated area to livestock including poultry.

### NOTICE TO BUYER

Greenshed Pty Ltd. will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.

PRIMO MAXX is a trademark of a Syngenta Group Company.

**APVMA APPROVAL NO.: 63628/58412**

**Batch No.:**

**Date of Manufacture:**

### FIRST AID

If swallowed DO NOT induce vomiting. Give a glass of water. If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (**Phone Australia 13 11 26**) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

### SAFETY DATA SHEET

Further information on this product is contained in a Safety Data Sheet (SDS) available from product suppliers or the manufacturer.

